

## Homework Assignment #10

## Note

This assignment is due 2:10PM Monday, May 27, 2019. Please write or type your answers on A4 (or similar size) paper. Put it on the instructor's desk before the class on the due date starts. Late submission will be penalized by 20% for each working day overdue. You may discuss the problems with others, but copying answers is strictly forbidden.

## Problems

(Note: problems marked with "Exercise X.XX" or "Problem X.XX" are taken from [Sipser 2013] with probable adaptation.)

- (Problem 5.9; 10 points) Let  $AMBIG_{CFG} = \{\langle G \rangle \mid G \text{ is an ambiguous CFG}\}$ . Show that  $AMBIG_{CFG}$  is undecidable. (Hint: use a reduction from PCP. Given an instance

$$P = \left\{ \left[ \frac{t_1}{b_1} \right], \left[ \frac{t_2}{b_2} \right], \dots, \left[ \frac{t_k}{b_k} \right] \right\}$$

of PCP, construct a CFG  $G$  with the rules:

$$\begin{aligned} S &\rightarrow T \mid B \\ T &\rightarrow t_1 T a_1 \mid \dots \mid t_k T a_k \mid t_1 a_1 \mid \dots \mid t_k a_k \\ B &\rightarrow t_1 B a_1 \mid \dots \mid t_k B a_k \mid t_1 a_1 \mid \dots \mid t_k a_k, \end{aligned}$$

where  $a_1, \dots, a_k$  are new terminal symbols. Prove that this reduction works.)

- (Problem 5.14(b); 20 points) Define a *two-headed finite automaton* (2DFA) to be a deterministic finite automaton that has two read-only, bidirectional heads that start at the left-hand end of the input tape and can be independently controlled to move in either direction. The tape of a 2DFA is finite and is just large enough to contain the input plus two additional blank tape cells, one on the left-end and one on the right-hand end, that serve as delimiters. A 2DFA accepts its input by entering a special accept state. For example, a 2DFA can recognize the language  $\{a^n b^n c^n \mid n \geq 0\}$ .

Let  $E_{2DFA} = \{\langle M \rangle \mid M \text{ is a 2DFA and } L(M) = \emptyset\}$ . Show that  $E_{2DFA}$  is undecidable.

- (Problem 5.18(c); 10 points) Use Rice's theorem to prove the undecidability of the language  $\{\langle M \rangle \mid M \text{ is a TM and } L(M) = \Sigma^*\}$ .
- (Problem 5.22; 20 points) Let  $X = \{\langle M, w \rangle \mid M \text{ is a single-tape TM that never modifies the portion of the tape that contains the input } w\}$ . Is  $X$  decidable? Prove your answer.
- (Problem 5.34; 20 points) Show that the Post Correspondence Problem is undecidable over the binary alphabet  $\Sigma = \{0, 1\}$ .
- (Problem 5.36; 20 points) Prove that there exists an undecidable subset of  $\{1\}^*$ .